

Records of *Beltrania rhombica* and *Didymobotryum rigidum* from China

XIAO-MEI WANG^{1*}, GUO-NING LIU², SHAN-SHAN CHEN¹ & XIAO-FENG DU¹

¹ College of Agronomy, Jilin Agricultural University, Changchun, 130118, China

² Baicheng Normal University, Baicheng, 137000, China

* CORRESPONDENCE TO: wxm820@126.com; communicate2007@126.com

ABSTRACT—*Beltrania rhombica* and *Didymobotryum rigidum* are described and illustrated, based on Chinese collections on decaying twigs of unidentified broadleaf plants. *Beltrania rhombica* represents a new record for mainland China, and *D. rigidum* a new record for Guangdong Province.

KEY WORDS—anamorphic fungi, taxonomy

Introduction

Beltrania Penz. is typified by *B. rhombica*, which was described from *Citrus limonum* [= *Citrus limon*] in Italy (Penzig 1882). This genus was described with distinguishing characters such as setae with radially lobed basal cells, conidiophores with separating cells, and biconic conidia with a hyaline transverse band and apical tubular appendage. *Didymobotryum* Sacc., which was established with *Periconia rigida* Berk. & Broome [≡ *D. rigidum*] as the type species, was mainly characterized by synnematos conidiophores and monotretic conidiogenous cells that produce unbranched acropetal chains of 1-septate conidia (Saccardo 1886).

Tropical and subtropical forest ecosystems offer particularly suitable habitats for saprobic microfungi, among which the anamorphic are the most abundant and diverse (e.g., Ma 2016; Ma et al. 2016; Xia et al. 2015a,b, 2016a,b). During a mycological survey of conidial fungi occurring on decaying plant material from Henan and Guangdong provinces, China, specimens of *Beltrania rhombica* and

Didymobotryum rigidum were collected on decaying twigs. The specimens are described and illustrated below and are deposited in the Herbarium of Mycology, Jilin Agricultural University, Changchun, China (HMJAU).

Beltrania rhombica Penz., Michelia 2(8): 474. 1882.

FIG. 1

CONIDIOPHORES macronematous, mononematous, arising in groups from the terminal cells of the immersed mycelium, erect, ascending, straight or



FIG. 1. *Beltrania rhombica* (HMJAU H6692). Conidiophores, conidiogenous cells, and conidia.

flexuous, unbranched, thick-walled, 4–8-septate, pale brown to mid brown, geniculate, conidial scars visible after secession of the conidia, $90\text{--}170 \times 3\text{--}5 \mu\text{m}$. CONIDIOGENOUS CELLS polyblastic, integrated, terminal, becoming lateral, determinate, cylindrical. CONIDIA solitary, dry, acrogenous or acropleurogenous, simple, biconic, smooth, slightly truncate at the base, with a rostrum at the apex, subhyaline to pale brown, with a hyaline median pale transverse band, $22\text{--}28 \times 7.5\text{--}10 \mu\text{m}$ (rostrum exclude), rostrum $4.5\text{--}7 \mu\text{m}$ long.

SPECIMEN EXAMINED: CHINA, HENAN PROVINCE, Jigong Mountain, on decaying twigs of an unidentified broadleaf tree, 2 Nov. 2016, J.W. Xia (HMJAU H6692).

COMMENTS—*Beltrania rhombica* is easily recognizable by its biconic conidia with a rostrum. Our specimen fits well with the original description of *B. rhombica* by Penzig (1882) except for the lack of setae. This species is reported for the first time from mainland China.

Didymobotryum rigidum (Berk. & Broome) Sacc., Syll. Fung. 4: 627. 1886. FIG. 2

CONIDIOMATA synnematal, unbranched, erect, with brown to dark brown stalks, consisting of compact aggregation of parallel conidiophores, terminating in brown fertile heads, $\leq 1000 \mu\text{m}$ long and $35\text{--}45 \mu\text{m}$ diam. in the middle, $\leq 110 \mu\text{m}$ diam. at the base, and $260 \mu\text{m}$ diam. at the apex. CONIDIOPHORES macronematous, erect, unbranched, septate, pale brown to brown, $2\text{--}4 \mu\text{m}$ diam. CONIDIOGENOUS CELLS monotretic, integrated, terminal, thick-walled, smooth, brown. Conidial secession schizolytic. CONIDIA solitary, dry, ellipsoidal, 1-euseptate, smooth, brown, $11\text{--}12.5 \times 4.5\text{--}6 \mu\text{m}$; septa $1.5\text{--}2.5 \mu\text{m}$ thick, darkly pigmented.

SPECIMEN EXAMINED: CHINA, GUANGDONG PROVINCE, Baiyun Mountain, on decaying twigs of an unidentified broadleaf tree, 6 Jun. 2015, J.W. Xia (HMJAU H6383).

COMMENTS—*Didymobotryum rigidum* is easily recognizable by its synnematous conidiophores and monotretic conidiogenous cells that produce catenate ellipsoidal conidia. This species has previously been reported from Sri Lanka, Thailand, Vietnam, China (Hong Kong), Japan, and Cuba (Berkeley & Broome 1873; Ellis 1971; Jones et al. 2004; Kobayashi 2007; Lu et al. 2000; Mercado Sierra et al. 1997; Minter et al. 2001; Saccardo 1886; Zhuang 2001); it is reported here for the first time from Guangdong Province.

Acknowledgments

The authors express gratitude to Dr. Xiu-Guo Zhang and Dr. Jian Ma for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (30600003). We acknowledge the facilities provided by Dr. P.M. Kirk and Drs. V. Robert, G. Stegehuis




FIG. 2. *Didymobotryum rigidum* (HMJAU H6383).
A. Synnema; B. Conidia; C–E. Conidiophores, conidiogenous cells, and conidia.

and Arthur Decock through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

Literature cited

- Berkeley MJ, Broome CE. 1873 ["1875"]. Enumeration of the fungi of Ceylon. Part II, containing the remainder of the hymenomycetes, with the remaining established tribes of fungi. Journal of the Linnean Society, Botany 14(74): 29–140.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, UK. 608 p.
- Jones EBG, Tanticharoen M, Hyde KD. 2004. Thai fungal diversity. BIOTEC, Pathum Thani, Thailand. 281 p.
- Kobayashi T. 2007. Index of fungi inhabiting woody plants in Japan: host, distribution and literature. Zenkoku-Nouson-Kyouiku-Kyoukai, Tokyo, Japan. 1227 p.
- Lu BS, Hyde KD, Ho WH, Tsui KM, Taylor JE, Wong KM, Yanna, Zhou DQ. 2000. Checklist of Hong Kong fungi. Fungal Diversity Research Series 5. 207 p.
- Ma J. 2016. *Corynesporopsis obclavata* and *Stanjehughesia jiangxiensis* spp. nov. from Lushan Mountain, China. Mycotaxon 131(3): 583–588. <https://doi.org/10.5248/131.583>
- Ma J, Zhang K, Zhang XG, Castañeda-Ruiz RE. 2016. Three new species of *Spadicoides* from Lushan Mountain, China. Mycological Progress 15: 43 [8 p.]. <https://doi.org/10.1007/s11557-016-1185-9>
- Mercado Sierra A, Holubová-Jechová V, Mena Portales J. 1997. Hifomicetes demaciáceos de Cuba: enteroblásticos. Museo Regionale di Scienze Naturali, Torino, Monografie 23. 388 p.
- Minter DW, Rodríguez Hernández M, Mena Portales J. 2001. Fungi of the Caribbean: an annotated checklist. PDMS Publishing Isleworth, UK. 946 p.
- Penzig O. 1882. Funghi agrumicoli: contribuzione allo studio dei funghi parassiti degli agrumi. Michelia 2(8): 385–508.
- Saccardo PA. 1886. Sylloge Hyphomycetum. Sylloge Fungorum 4. 807 p.
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG. 2015a. *Sporidesmiopsis malloti* sp. nov. and new records from southern China. Mycotaxon 130(3): 827–833. <https://doi.org/10.5248/130.827>
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG. 2015b. *Codinaea jianfenglingensis* sp. nov. and new records from southern China. Mycotaxon 130(3): 835–841. <https://doi.org/10.5248/130.835>
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG. 2016a. *Sympodiosynnema*, a new genus of dematiaceous hyphomycetes from southern China. Mycotaxon 131(1): 45–48. <https://doi.org/10.5248/131.45>
- Xia JW, Ma YR, Gao JM, Zhang XG, Li Z. 2016b. Two new species of *Endophragmiella* from southern China. Nova Hedwigia 103: 349–357. https://doi.org/10.1127/nova_hedwigia/2016/0356
- Zhuang WY. 2001. Higher fungi of tropical China. Mycotaxon Ltd., Ithaca NY, USA. 485 p.